

PEX5, the Shuttling Import Receptor for Peroxisomal Matrix Proteins, Is a Redox-Sensitive Protein

Oksana Apanasets¹, Cláudia P. Grou², Paul P. Van Veldhoven¹, Chantal Brees¹, Bo Wang¹, Marcus Nordgren¹, Gabriele Dodt³, Jorge E. Azevedo^{2,4}, Marc Fransen^{1,*}

This is the peer reviewed version of the following article: Apanasets, O., Grou, C. P., Van Veldhoven, P. P., Brees, C., Wang, B., Nordgren, M., Dodt, G., Azevedo, J. E. and Fransen, M. (2014), PEX5, the Shuttling Import Receptor for Peroxisomal Matrix Proteins, Is a Redox - Sensitive Protein. *Traffic*, 15: 94-103., which has been published in final form at <https://doi.org/10.1111/tra.12129>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.

¹Laboratory of Lipid Biochemistry and Protein Interactions, Department of Cellular and Molecular Medicine, KU, Leuven, Belgium

²Instituto de Biologia Molecular e Celular (IBMC), Universidade do Porto, Porto, Portugal

³Interfakultäres Institut für Biochemie, UniversitätTübingen, Tübingen, Germany

⁴Instituto de Ciências Biomédicas Abel Salazar (ICBAS), Universidade do Porto, Porto, Portugal

*Corresponding author: Marc Fransen, marc.fransen@med.kuleuven.be

ABSTRACT

Peroxisome maintenance depends on the import of nuclear-encoded proteins from the cytosol. The vast majority of these proteins is destined for the peroxisomal lumen and contains a C-terminal peroxisomal targeting signal, called PTS₁. This targeting signal is recognized in the cytosol by the receptor PEX5. After docking at the peroxisomal membrane and release of the cargo into the organelle matrix, PEX5 is recycled to the cytosol through a process requiring monoubiquitination of an N-terminal, cytosolically exposed cysteine residue (Cys11 in the human protein). At present, the reason why a cysteine, and not a lysine residue, is the target of ubiquitination remains unclear. Here, we provide evidence that PTS₁ protein import into human fibroblasts is a redox-sensitive process. We also demonstrate that Cys11 in human PEX5 functions as a redox switch that regulates PEX5 activity in response to intracellular oxidative stress. Finally, we show that exposure of human

INSTITUTO
DE INVESTIGAÇÃO
E INOVAÇÃO
EM SAÚDE
UNIVERSIDADE
DO PORTO

Rua Alfredo Allen, 208
4200-135 Porto
Portugal
+351 220 408 800
info@i3s.up.pt
www.i3s.up.pt

Version: Postprint (identical content as published paper) This is a self-archived document from i3S – Instituto de Investigação e Inovação em Saúde in the University of Porto Open Repository For Open Access to more of our publications, please visit <http://repositorio-aberto.up.pt/>

PEX5 to oxidized glutathione results in a ubiquitination-deficient PEX5 molecule, and that substitution of Cys11 by a lysine can counteract this effect. In summary, these findings reveal that the activity of PEX5, and hence PTS1 import, is controlled by the redox state of the cytosol. The potential physiological implications of these findings are discussed.

INTRODUCTION

Peroxisomes are multifunctional organelles that, among other functions, participate in cellular lipid and reactive oxygen species (ROS) metabolism (1,2). Defects in peroxisome function are at the core of several fatal genetic disorders (3). In addition, there is growing evidence that peroxisome biology and cellular aging are closely intertwined (4). For example, inhibition of the activity of catalase, the most abundant peroxisomal antioxidant enzyme, causes a cascade of accelerated aging reactions such as increased mitochondrial ROS production, oxidative damage to proteins and DNA, hypersecretion of matrix metalloproteinases, and impaired cell growth (5,6). On the other hand, increased cellular oxidative stress (e.g. during cellular aging) leads to a reduced import efficiency of peroxisomal matrix proteins containing a C-terminal peroxisomal targeting signal type 1 (PTS1), especially catalase (7). The molecular mechanisms underlying these phenomena are far from being understood. However, in this context, it should be mentioned that (i) under conditions of oxidative stress, PEX5, the shuttling import receptor for PTS1 proteins, accumulates at the peroxisome membrane and (ii) the cellular senescence phenotypes in old cells can be reversed by overexpression of catalase-SKL, a catalase derivative with enhanced peroxisome targeting efficiency (8).

The assembly of functional peroxisomes requires the import of both membrane and matrix proteins (9). The vast majority of matrix proteins contain a PTS1. This class of proteins is synthesized on free polyribosomes in the cytosol and post-translationally recognized by the shuttling import receptor PEX5 while still in the cytosol. The PEX5- cargo protein complex then interacts with the peroxisomal docking/translocation machinery (DTM). This interaction ultimately leads to the insertion of PEX5 into the DTM with the concomitant translocation of the cargo protein across the peroxisomal membrane. Interestingly, a 2-kDa N-terminal region of PEX5 containing a phylogenetically conserved cysteine residue remains exposed to the cytosol during this process. After cargo release, PEX5 is monoubiquitinated at this conserved cysteine residue (Cys11), and extracted from the DTM by the ATP-dependent receptor export module (REM) (10–12). Once back in the cytosol, the ubiquitin moiety attached to PEX5 is removed, probably by a combination of enzymatic and non-enzymatic mechanisms (13,14). This resets PEX5 for the next round of import.

Protein ubiquitination is a posttranslational modification that controls many intracellular processes (15,16). In general, this modification results in the attachment of a ubiquitin molecule to an ϵ -amino group of a substrate lysine residue (17). However, in a select set of proteins (e.g. PEX5), ubiquitin is transferred onto a cysteine residue yielding a thioester (18). The physiological relevance and role of this unconventional type of ubiquitination are puzzling. On one hand, it is well known that thioesters are easily disrupted by nucleophiles such as reduced glutathione (GSH) (19). On the other hand, it has been reported that substitution of the cysteine at position 11 by lysine in human PEX5 results in a fully functional PTS1 import receptor, at least under standard laboratory conditions (13). Prompted by the combined observations that (i) PEX5 accumulates at the peroxisomal membrane upon oxidative stress (8), (ii) this peroxin needs to be monoubiquitinated at a conserved cysteine

residue in order to be exported back to the cytosol (20,21) and (iii) there is an ever increasing number of examples in which the activity of proteins (e.g. transcription factors, kinases, phosphatases, etc.) is controlled by the oxidation states of cysteines (22), we recently hypothesized that Cys11 in human PEX5 may serve as a functional redox switch regulating the peroxisomal/cytosolic localization of peroxisomal proteins such as catalase (2). In this report, we provide experimental evidence in favor of this hypothesis.

Materials and Methods

DNA manipulations and plasmids

The mammalian expression vector pIRES2-EGFP and the restriction enzymes were commercially obtained from Westburg. Oligonucleotides (Table S1, Supporting Information) were synthesized by Integrated DNA Technologies. Polymerase chain reactions were routinely carried out by using Pfx DNA Polymerase (Invitrogen). The *Escherichia coli* strain Top10F' (Invitrogen) was used for all DNA manipulations. The plasmids encoding a cytosolic roGFP2 (roGFP2; pMF1707), peroxisomal roGFP2 (roGFP2-PTS1; pMF1706) or human catalase (pMF1525) have been described elsewhere (23). The pIRES2-roGFP2-PTS1 (pMN8) plasmid was constructed via a three-point ligation of the following fragments: (i) the XmaI/NotI-restricted backbone fragment of pIRES2-EGFP, (ii) a XmaI/BamHI-restricted polymerase chain reaction (PCR) fragment comprising the IRES2 sequence (template: pIRES2-EGFP2; primers: pIRES2fwXmaI and pIRES2rvBamHI) and (iii) the BamHI/NotI restricted fragment of pMF1706, which encodes roGFP2-PTS1. The bicistronic constructs encoding roGFP2-PTS1 in combination with the large isoform of HsPEX5WT (pDC2) or HsPEX5C11K (pDC1) were generated by amplifying the corresponding HsPEX5-encoding cDNA fragments by PCR (templates: pMF1678 and pMF1578, respectively (13); primers: HsPex5.1fw and HsPex5.2rv) and cloning the BglII/Sall-digested PCR products into the BglII/Sall-restricted pIRES2-roGFP2-PTS1 vector. All plasmids were verified by DNA sequencing (LGC Genomics). The pGEM4(Promega)-based plasmids encoding the large isoform of human PEX5 and the mutated version of it possessing a lysine at position 11 (PEX5C11K) have been previously described (13,43).

Cell culture, transfections and fluorescence microscopy

The primary PEX5 null HuFs (cell line PBD005) have been described elsewhere (44,45). Briefly, these cells are homozygous for a mutation (Arg390Ter) in the PEX5 gene that results in a complete inactivation of PEX5's normal function (44,45). Control HuFs were kindly provided by Dr. D. Cassiman (KU, Leuven, Belgium). Where indicated, the cell lines were transformed by introduction of SV40 large T-antigen. Unless specified otherwise, all cell lines were cultivated at 37° C in a humidified 5% CO₂ incubator in minimum essential medium Eagle alpha (MEMα; Lonza) supplemented with 10% (v/v) heat-inactivated fetal bovine serum (Invitrogen), 2 mM Glutamax (Invitrogen), and MycoZap (Lonza). The cells were transfected by using the Neon Transfection System (Invitrogen; primary HuFs: 1200 V, 20-milliseconds pulse width, 1 pulse; transformed HuFs: 1300 V, 30-milliseconds pulse width, 1 pulse). Cells for live cell imaging were seeded and imaged in FD-35 Fluorodish cell culture dishes (World Precision Instruments). The technical specifications of the objectives, excitation and emission filters, and digital camera have been described elsewhere (46). Ratiometric redox measurements were performed as described (23). Note that, as the spectral output of the replacement lamps was not always identical, the emission ratios of roGFP2 at the 400-

and 480-nm excitation wavelengths were expressed in arbitrary units (a.u.). The Olympus image analysis and particle detection software were used for quantitative image analysis.

Peroxisome-dependent PEX5-ubiquitination assay

³⁵S-labeled PEX5 and PEX5C11K were synthesized in rabbit reticulocyte lysates using the TNT®quick coupled transcription/translation system (Promega) in the presence of EasyTag™ L-(³⁵S)methionine (specific activity, >1000 Ci/mmol; PerkinElmer Life Sciences) and the pGEM4-based plasmids referred to above. Rat liver PNS (600 µg of protein) was diluted in import buffer (0.25 M sucrose, 50 mM KCl, 20 mM MOPS-KOH pH 7.2, 3 mM MgCl₂, 20 µM methionine, and 2 µg/mL N-(trans-epoxysuccinyl)-L-leucine 4-guanidinobutylamide) containing 0.3 mM ATP (final volume 80 µL), and primed for import by incubating the samples for 5 min at 37° C (47). The samples were placed on ice, and incubated for 10 min in the absence or presence of increasing concentrations oxidized glutathione (GSSG; added from a 0.1 M stock solution in 20 mM MOPS-KOH, pH 7.4). In parallel, 1 µL of ³⁵S-labeled PEX5WT or PEX5C11K (diluted 1:20 (v/v) in import buffer) were also treated with the matching GSSG concentration. Next, the diluted lysates were added to the corresponding PNS mixtures, and the samples were incubated for 20 min at 37° C in the presence of 5 mM ATP and 7 µM glutathione S-transferase (GST)-ubiquitin (20). The reactions were stopped with 20 mM N-ethylmaleimide (NEM; 5 min on ice), diluted with 900 µL of ice-cold SEMK buffer (0.25 M sucrose, 80 mM KCl, 20 mM MOPS-KOH, pH 7.2, 1 mM EDTA-NaOH, pH 8.0), and centrifuged at 16 000 × g for 20 min at 4° C to separate the organelles from soluble proteins. The organelle pellets were then subjected to SDS-PAGE under non-reducing (no DTT) and reducing (0.1 M DTT) conditions, and analyzed by Western blot and autoradiography, as described before (13,20).

Statistical analysis

Statistics were performed on the VassarStats statistical computation website (<http://vassarStats.net/>). One-way analysis of variance was used to determine the differences among independent groups of numerical values, and individual differences were further explored with a Student's t-test. The significance level was chosen to be $p < 0.01$.

Results

Redox regulation of PTS1 protein import into peroxisomes

Previously, we and others have shown that there is a strong correlation between cellular aging, oxidative stress and PTS1 protein import into peroxisomes (7,8,23). To strengthen the concept that changes in the cellular redox state may have wide-ranging consequences for peroxisomal matrix protein import, we cultivated human fibroblasts (HuFs; passage 35) under different redox conditions and studied the subcellular localization of roGFP2-PTS1, a peroxisome-targeted variant of roGFP2 (Figure 1A). RoGFP2 is a redox-sensitive form of the enhanced green fluorescent protein (EGFP) that contains two engineered cysteine residues on adjacent surface-exposed strands close to the chromophore (24). As (i) roGFP2 has two fluorescence excitation maxima at ~400 and ~490 nm and (ii) disulfide bond formation between the engineered cysteine residues leads to an increase in the excitation peak at ~400 nm at the expense of the ~490 nm peak, the ratio of roGFP2 emissions (at ~510 nm) can provide a non-destructive read-out of the redox environment of the chromophore (24). To determine the effects of different treatments on the redox state of the cytosol, we

employed non-targeted, i.e. cytosolic, roGFP2 (23). We found that overexpression of catalase, a peroxisomal H₂O₂-decomposing enzyme whose activity drops during in vitro cellular aging (25), significantly lowers the redox state of the cytosol (Figure 1B) and largely restores PTS1 protein import (Figure 1C). A similar result was obtained when the cells were cultivated overnight in medium containing 0.5 mM dithiothreitol (DTT), a membrane permeable reducing agent (Figure 1B,C). For comparison, we also included the values we obtained for these fibroblasts at passage 15 (Figure 1B,C). Taken together, these findings lend support to the idea that the import of PTS1 proteins into peroxisomes is a redox-regulated process.

The redox-dependent changes in PTS1 import efficiency involve the evolutionarily conserved cysteine residue in the N-terminal domain of PEX5

As (i) PTS1 proteins are imported into peroxisomes by the shuttling receptor PEX5 (9), (ii) this receptor accumulates at the peroxisomal membrane in aging cells (7), and (iii) the export of PEX5 from the peroxisomal membrane to the cytosol depends on the evolutionarily conserved cysteine residue in the N-terminal region of the protein (20), we investigated whether or not this cysteine can act as a redox-sensitive regulatory switch for PEX5 activity. Therefore, we complemented late-passage primary HuFs lacking functional PEX5 (hereafter called PEX5 null; for more details, see Materials and Methods) with constructs encoding human PEX5WT or PEX5C11K and determined how efficiently the expression of both PEX5 variants restored PTS1 protein import. The reasoning behind this experiment was that – in case Cys11 does indeed function as a redox switch – PEX5C11K, a peroxisomal import/export-competent protein (13), should restore PTS1 import more efficiently than PEX5WT in cells with an increased cytosolic redox state. To exclude the possibility that any observed differences are the indirect result of non-redox-regulated events, we also performed similar experiments in the corresponding transformed HuFs, which display a lower cytosolic redox state (see below). The results of these experiments are presented in Figure 2A. From these data, it is clear that (i) the import of roGFP2-PTS1 into peroxisomes is PEX5-dependent, (ii) PEX5WT is less functional in late-passage primary cells than in transformed cells and (iii) the replacement of cysteine at position 11 by a lysine in PEX5 dramatically increases its functionality in late-passage primary cells. To provide unambiguous evidence that the differences observed in primary and transformed cells expressing PEX5WT and PEX5C11K are redox-related and not simply due to differences in expression or stability of these PEX5 proteins, we confirmed that transformation of the PEX5 null HuFs led to a more reduced redox state of the cytosol (Figure 2B) and studied the effect of the thiol-reducing agent N-acetyl-cysteine (NAC) on the complementation activity of PEX5WT and PEX5C11K (Figure 3). Importantly, treatment of the cells with NAC, a compound that shifts the redox state of the cytosol toward lower values (26) (Figure 3A), resulted in an increased functionality of PEX5WT, but not of PEX5C11K (Figure 3B,C). Note that the results obtained for PEX5C11K also exclude the possibility that the differences in import kinetics observed for PEX5WT upon NAC treatment are the result of redox-related structural changes of the roGFP2-PTS1 reporter protein. All together, these experiments suggest that Cys11 of human PEX5 functions as a redox switch that modulates PEX5 activity in response to intracellular oxidative stress.

Exposure of human PEX5 to oxidized glutathione impairs its monoubiquitination in a Cys11-dependent manner

To determine whether or not oxidation of human PEX5 affects its monoubiquitination at Cys11, we incubated ³⁵S-labeled PEX5WT or PEX5C11K with a (peroxisome-containing) rat liver post-nuclear

supernatant (PNS) supplemented with ATP and GST-tagged ubiquitin (GST-Ub) in the absence or presence of increasing concentrations of oxidized glutathione (GSSG; for detailed procedures, see Materials and Methods). Organelles were then processed for SDS–PAGE under non-reducing and reducing conditions, and the amounts of ³⁵S-labeled GST-Ub-PEX₅ species that were produced in these in vitro assays were assessed by autoradiography (Figure 4). As shown in Figure 4 (upper panel), monoubiquitination of PEX₅WT is much more sensitive to GSSG than monoubiquitination of PEX₅C11K. Note that PEX₅ monoubiquitination detected in these assays occurs at its residue 11, because treatment of protein samples with DTT destroys the thioester bond between GST-Ub and PEX₅WT but not the amide bond that links GST-Ub to PEX₅C11K (Figure 4, lower panel; see also 13). The reduction in PEX₅C11K monoubiquitination observed with the highest concentration of GSSG (Figure 4, upper panel) is probably due to the fact that the ubiquitination cascade comprises cysteine-containing enzymes (the ubiquitin-activating enzyme and ubiquitin-conjugating enzymes) that are also potential targets of GSSG (27). In summary, these data strongly indicate that Cys11 in human PEX₅ is the most oxidative stress-sensitive cysteine of the complete PEX₅ monoubiquitination process.

Discussion

In this study, we have provided experimental evidence that (i) PTS₁ protein import is a redox-sensitive process, (ii) this redox sensitivity is mediated by PEX₅, the shuttling receptor for peroxisomal matrix proteins, and (iii) the cysteine at position 11 in human PEX₅ functions as a redox switch that regulates PEX₅ activity in response to oxidative stress, most likely by modulating its monoubiquitination at the DTM. In summary, these findings provide a molecular explanation for the long-standing observation that peroxisomal PTS₁ protein import is reduced in cells experiencing oxidative stress (7,28). In addition, as PEX₅ monoubiquitination at Cys11 is necessary for the export of the receptor back into the cytosol (20,21), these data also explain why PEX₅ accumulates on peroxisomal membranes during cellular aging (7), a condition associated with a sharp increase in the cytosolic redox state (23,29). In the context of these results, it is important to recognize that also additional mechanisms may be at play in relation to the inhibitory effects of oxidative stress on peroxisomal protein import. For example, as oxidative stress can decrease both glycolytic and mitochondrial ATP production (30,31), it is very likely that this type of stress can also impede the ATP-dependent export step of PEX₅ from the peroxisomal membrane to the cytosol. In addition, one can easily envisage a scenario in which oxidative stress leads to the modification and partial inactivation of non-PEX₅-related components of the peroxisomal protein import and protein quality control machineries (32,33). However, as (i) such alternative mechanisms can be expected to equally reduce the functionality of PEX₅WT and PEX₅C11K and (ii) this is not the case in the experiments discussed here, they seem to be of no importance under the conditions studied.

Over the years, accumulating evidence has strengthened the view that peroxisomal metabolism, cellular oxidative balance and cellular aging are intimately interconnected (2,4,34). In the context of this work, it is important to remember that Terlecky and colleagues in 2006 postulated the existence of a self-sustaining peroxisome deterioration spiral (35). According to this model, peroxisomes are predisposed to oxidative stress due to a dysregulated import of peroxisomal H₂O₂-producing oxidases, which in general bear a strong PTS₁ (36), and catalase, which contains a weak PTS₁ (37). This causes the organelle to slowly lose its ability to clear generated ROS, and – as such – its import machinery may become gradually compromised. This, in turn, will initially mainly

affect the import of weak substrates, such as catalase, but later on result in a self-perpetuating negative protein import spiral causing oxidative stress and peroxisome dysfunction (6,35). The data presented in this manuscript extend this theory by providing specific details regarding the potential mechanisms by which cellular oxidative stress causes peroxisomal protein import defects. Note that, as the monoubiquitinable cysteine residue in the N-terminus of PEX5 and the weak PTS1 of catalase are highly conserved in virtually all mammalian species (data not shown), it is tempting to speculate that these features have evolved because of a possible advantage. For example, a decrease in the peroxisomal import efficiency of (newly-synthesized) catalase may allow cells to rapidly respond to oxidative stress in the cytosol. This postulate is in line with a recent study showing that p97, an ATPase of the AAA family (ATPases with diverse cellular activities), can regulate H₂O₂ levels in the cytosol by affecting the retention time of (newly-synthesized) catalase within this subcellular compartment (38).

Importantly, when the work presented in this manuscript was in a final stage, Subramani and coworkers reported that also *Pichia pastoris* Pex5 (PpPex5) functions as a redox-regulated PTS1 protein import receptor (39). Specifically, these authors reported that Cys10, the evolutionarily conserved redox-sensitive cysteine in PpPex5, acts as a redox switch to control the binding and release of PTS1 proteins to and from PpPex5 by modulating its oligomeric state. On the basis of these observations and a report stating that the intraperoxisomal redox state is more reductive than that of the cytosol (40), the authors postulated a working model in which (oxidized) disulfide bond-linked PpPex5 binds newly-synthesized PTS1 proteins in the cytosol and releases these proteins upon reduction in the peroxisomal matrix. However, here it is important to note that this model is not compatible with our data. Indeed, we clearly show that the reduced functionality of PEX5 in an oxidizing environment can be counteracted by substituting Cys11 with lysine, and this finding in turn strongly indicates that – at least for human PEX5 – oxidation-mediated disulfide bond formation through this conserved cysteine can be functionally uncoupled from PTS1 cargo binding and release. In this context, it is also important to point out that some of us have previously shown that, in a mammalian system, (i) PTS1 proteins can be efficiently imported into peroxisomes under conditions in which the redox state of peroxisomes is more oxidizing than that of the cytosol (23), (ii) the release of PTS1 cargo proteins into the peroxisomal matrix is most likely triggered by binding of PEX14 to the 6th (and probably 7th) di-aromatic motif(s) of PEX5 (41) and (iii) PEX5C11A, a mutant of PEX5 that is not capable of forming a disulfide bond through the conserved cysteine, can translocate PTS1 proteins across the peroxisomal membrane and release these proteins into the organellar matrix as efficient as PEX5WT, at least in a cell-free in vitro import system (as PEX5C11A cannot be monoubiquitinated at the peroxisomal membrane and hence not be exported back to the cytosol, this protein abrogates PTS1 protein import upon expression in wild-type cells) (21,42). Nevertheless, as yeast and mammalian systems are not always directly comparable because of differences in PEX proteins and enzyme content (9), it remains to be established whether or not the apparently conflicting data obtained for PpPex5p and human PEX5 represent species- or condition-specific characteristics. In conclusion, our data provide clear evidence that (i) the activity of PEX5 depends on the redox state of the cytosol and (ii) this redox sensitivity is mediated by Cys11, an evolutionarily conserved amino acid involved in receptor monoubiquitination and recycling. These findings are relevant for a better understanding of how cellular oxidative stress may contribute to a demise of peroxisome function, a condition associated with cellular senescence, organismal aging and age-related diseases (4).

Acknowledgements

The authors are grateful to Dr. Ann Moser (Baltimore, USA) for the primary PEX5 null human fibroblasts. This work was supported by grants from the 'Fonds voor Wetenschappelijk Onderzoek-Vlaanderen (Onderzoeksproject G.0754.09)' (to M. F. and P. P. V. V.), by the KU Leuven grants OT/09/045 (to M. F. and P. P. V. V.) and DBOF/10/059 (to P. P. V. V. and M. F.), and by FEDER funds through the Operational Competitiveness Programme – COMPETE and by National Funds through FCT – Fundação para a Ciência e a Tecnologia under the project FCOMP-01-0124-FEDER-019731 (PTDC/BIA-BCM/118577/2010) (to J. E. A.). M. N. is supported by a FLOF fellowship from the Department of Cellular and Molecular Medicine, KU Leuven. B. W. is a recipient of a DBOF fellowship (DBOF/10/059) from the KU Leuven. C. P. G. is supported by Fundação para a Ciência e a Tecnologia, Programa Operacional Potencial Humano do QREN, and Fundo Social Europeu.

REFERENCES

1. Van Veldhoven PP. Biochemistry and genetics of inherited disorders of peroxisomal fatty acid metabolism. *J Lipid Res* 2010;51:2863–2895.
2. Fransen M, Nordgren M, Wang B, Apanasets O. Role of peroxisomes in ROS/RNS-metabolism: implications for human disease. *Biochim Biophys Acta* 1822;2012:1363–1373.
3. Waterham HR, Ebberink MS. Genetics and molecular basis of human peroxisome biogenesis disorders. *Biochim Biophys Acta* 1822;2012:1430–1441.
4. Fransen M, Nordgren M, Wang B, Apanasets O, Van Veldhoven PP. Aging, age-related diseases and peroxisomes. *Subcell Biochem* 2013;69:45–65.
5. Koepke JI, Wood CS, Terlecky LJ, Walton PA, Terlecky SR. Progeric effects of catalase inactivation in human cells. *Toxicol Appl Pharmacol* 2008;232:99–108.
6. Walton PA, Pizzitelli M. Effects of peroxisomal catalase inhibition on mitochondrial function. *Front Physiol* 2012;3:108.doi:10.3389/fphys.2012.00108.
7. Legakis JE, Koepke JI, Jedeszko C, Barlasakar F, Terlecky LJ, Edwards HJ, Walton PA, Terlecky SR. Peroxisome senescence in human fibroblasts. *Mol Biol Cell* 2002;13:4243–4255.
8. Koepke JI, Nakrieko KA, Wood CS, Boucher KK, Terlecky LJ, Walton PA, Terlecky SR. Restoration of peroxisomal catalase import in a model of human cellular aging. *Traffic* 2007;8:1590–1600.
9. Fransen M. Peroxisome dynamics: molecular players, mechanisms, and (dys)functions. *ISRN Cell Biol* 2012 Article ID 714192.
10. Grou CP, Carvalho AF, Pinto MP, Alencastre IS, Rodrigues TA, Freitas MO, Francisco T, Sá-Miranda C, Azevedo JE. The peroxisomal protein import machinery – a case report of transient ubiquitination with a new flavor. *Cell Mol Life Sci* 2009;66:254–262.

11. Ma C, Agrawal G, Subramani S. Peroxisome assembly: matrix and membrane protein biogenesis. *J Cell Biol* 2011;193:7–16.
12. Platta HW, Hagen S, Erdmann R. The exportomer: the peroxisomal receptor export machinery. *Cell Mol Life Sci* 2013;70:1393–1411.
13. Grou CP, Carvalho AF, Pinto MP, Huybrechts SJ, Sá-Miranda C, Fransen M, Azevedo JE. Properties of the ubiquitin-PEX5 thiol ester conjugate. *J Biol Chem* 2009;284:10504–10513.
14. Grou CP, Francisco T, Rodrigues TA, Freitas MO, Pinto MP, Carvalho AF, Domingues P, Wood SA, Rodríguez-Borges JE, Sá-Miranda C, Fransen M, Azevedo JE. Identification of ubiquitin-specific protease 9X (USP9X) as a deubiquitinase acting on ubiquitin-peroxin 5 (PEX5) thioester conjugate. *J Biol Chem* 2012;287:12815–12827.
15. Chen ZJ, Sun LJ. Nonproteolytic functions of ubiquitin in cell signaling. *Mol Cell* 2009;33:275–286.
16. Clague MJ, Urbe S. Ubiquitin: same molecule, different degradation pathways. *Cell* 2010;143:682–685.
17. Kommander D. The emerging complexity of protein ubiquitination. *Biochem Soc Trans* 2009;37:937–953.
18. Wang X, Herr RA, Hansen TH. Ubiquitination of substrates by esterification. *Traffic* 2012;13:19–24.
19. Rose IA, Warms JV. An enzyme with ubiquitin carboxy-terminal esterase activity from reticulocytes. *Biochemistry* 1983;22:4234–4237.
20. Carvalho AF, Pinto MP, Grou CP, Alencastre IS, Fransen M, Sá-Miranda C, Azevedo JE. Ubiquitination of mammalian PEX5, the peroxisomal import receptor. *J Biol Chem* 2007;282:31267–33172.
21. Okumoto K, Misono S, Miyata N, Matsumoto Y, Mukai S, Fujiki Y. Cysteine ubiquitination of PTS1 receptor PEX5 regulates PEX5 recycling. *Traffic* 2011;12:1067–1083.
22. Pace NJ, Weerapana E. Diverse functional roles of reactive cysteines. *ACS Chem Biol* 2013;8:283–296.
23. Ivashchenko O, Van Veldhoven PP, Brees C, Ho YS, Terlecky SR, Fransen M. Intraperoxisomal redox balance in mammalian cells: oxidative stress and interorganellar cross-talk. *Mol Biol Cell* 2011;22:1440–1451.
24. Hanson GT, Aggeler R, Oglesbee D, Cannon M, Capaldi RA, Tsien RY, Remington SJ. Investigating mitochondrial redox potential with redox-sensitive green fluorescent protein indicators. *J Biol Chem* 2004;279:13044–13053.

25. Kaneko T, Tahara S, Taguchi T, Kondo H. Accumulation of oxidative DNA damage, 8-oxo-2-deoxyguanosine, and change of repair systems during in vitro cellular aging of cultured human skin fibroblasts. *Mutat Res* 2001;487:19–30.
26. Wu G, Fang YZ, Yang S, Lupton JR, Turner ND. Glutathione metabolism and its implications for health. *J Nutr* 2004;134:489–492.
27. Jahngen-Hodge J, Obin MS, Gong X, Shang F, Nowell TR Jr, Gong J, Abasi H, Blumberg J, Taylor A. Regulation of ubiquitin-conjugating enzymes by glutathione following oxidative stress. *J Biol Chem* 1997;272:28218–28226.
28. Sheikh FG, Pahan K, Khan M, Barbosa E, Singh I. Abnormality in catalase import into peroxisomes leads to severe neurological disorder. *Proc Natl Acad Sci U S A* 1998;95:2961–2966.
29. Muller M. Cellular senescence: molecular mechanisms, in vivo significance, and redox considerations. *Antioxid Redox Signal* 2009;11:59–98.
30. Taha R, Seidman E, Mailhot G, Boudreau F, Gendron FP, Beaulieu JF, Menard D, Delvin E, Amre D, Levy E. Oxidative stress and mitochondrial functions in the intestinal Caco-2/15 cell line. *PLoS One* 2010;27:e11817. doi:10.1371/journal.pone.0011817.
31. Rovito HA, Oblong JE. Nicotinamide preferentially protects glycolysis in dermal fibroblasts under oxidative stress conditions. *Br J Dermatol* 2013;169:15–24.
32. Nordgren M, Fransen M. Peroxisomal metabolism and oxidative stress. *Biochimie* 2013 (In press). doi:10.1016/j.biochi.2013.07.026.
33. Omi S, Nakata R, Okamura-Ikeda K, Konishi H, Taniguchi H. Contribution of peroxisome-specific isoform of Lon protease in sorting PTS1 proteins to peroxisomes. *J Biochem* 2008;143:649–660.
34. Titorenko VI, Terlecky SR. Peroxisome metabolism and cellular aging. *Traffic* 2011;12:252–259.
35. Terlecky SR, Koepke JI, Walton PA. Peroxisomes and aging. *Biochim Biophys Acta* 2006;1763:1749–1754.
36. Lametschwandtnr G, Brocard C, Fransen M, Van Veldhoven P, Berger J, Hartig A. The difference in recognition of terminal tripeptides as peroxisomal targeting signal 1 between yeast and human is due to different affinities of their receptor PEX5 to the cognate signal and to residues adjacent to it. *J Biol Chem* 1998;273:33635–33643.
37. Purdue PE, Lazarow PB. Targeting of human catalase to peroxisomes is dependent upon a novel COOH-terminal peroxisomal targeting sequence. *J Cell Biol* 1996;134:849–862.
38. Murakami K, Ichinohe Y, Koike M, Sasaoka N, Iemura S, Natsume T, Kakizuka A. VCP is an integral component of a novel feedback mechanism that controls intracellular localization of catalase and H₂O₂ levels. *PLoS One* 2013;8:e56012.

39. Ma C, Hagstrom D, Polley SG, Subramani S. Redox regulated cargo binding and release by the peroxisomal targeting signal receptor, Pex5. *J Biol Chem* 2013;288:27220–27231. doi:10.1074/jbc.M113.492694.
40. Yano T, Oku M, Akeyama N, Itoyama A, Yurimoto H, Kuge S, Fujiki Y, Sakai Y. A novel fluorescent sensor protein for visualization of redox states in the cytoplasm and in peroxisomes. *Mol Cell Biol* 2010;30:3758–3766.
41. Freitas MO, Francisco T, Rodrigues TA, Alencastre IS, Pinto MP, Grou CP, Carvalho AF, Fransen M, Sá-Miranda C, Azevedo JE. PEX5 protein binds monomeric catalase blocking its tetramerization and releases it upon binding the N-terminal domain of PEX14. *J Biol Chem* 2011;286:40509–40519.
42. Francisco T, Rodrigues TA, Freitas MO, Grou CP, Carvalho AF, Sá-Miranda C, Pinto MP, Azevedo JE. A cargo-centered perspective on the PEX5-mediated peroxisomal protein import pathway. *J Biol Chem* 2013;288:29151–29159. doi:10.1074/jbc.M113.487140.
43. Gouveia AM, Guimaraes CP, Oliveira ME, Reguenga C, Sa-Miranda C, Azevedo JE. Characterization of the peroxisomal cycling receptor, Pex5p, using a cell-free in vitro import system. *J Biol Chem* 2003;278:226–232.
44. Dodt G, Braverman N, Wong C, Moser A, Moser HW, Watkins P, Valle D, Gould SJ. Mutations in the PTS1 receptor gene, PXR1, define complementation group 2 of the peroxisome biogenesis disorders. *Nat Genet* 1995;9:115–125.
45. Slawewski ML, Dodt G, Steinberg S, Moser AB, Moser HW, Gould SJ. Identification of three distinct peroxisomal protein import defects in patients with peroxisome biogenesis disorders. *J Cell Sci* 1995;108:1817–1829.
46. Nordgren M, Wang B, Apanasets O, Brees C, Veldhoven PP, Fransen M. Potential limitations in the use of KillerRed for fluorescence microscopy. *J Microsc* 2012;245:229–235.
47. Oliveira ME, Gouveia AM, Pinto RA, Sá-Miranda C, Azevedo JE. The energetics of Pex5p-mediated peroxisomal protein import. *J Biol Chem* 2003;278:39483–39488.




Figure 1: The efficiency of PTS1 protein import into peroxisomes is influenced by the cytosolic redox state. Primary HuFs (passages 15 or 35) were transiently transfected with a plasmid encoding peroxisomal (roGFP2-PTS1) or cytosolic (roGFP2) roGFP2, in combination or not with a plasmid encoding wild-type catalase (cat). The cells were cultivated in standard growth medium and, where indicated, supplemented with 0.5 mM DTT (overnight before analysis). A) Subcellular localizations of roGFP2-PTS1 in late passage cells. Note that, as (i) the cytosolic redox state of these cells displays a considerable intercellular variability (23), and (ii) this variability substantially influences the import efficiency of PTS1 proteins (this study), different cells can display a different distribution pattern for roGFP2-PTS1. PO, peroxisomes; C, cytosol; PO/C, peroxisomes and cytosol. Scale bar: 10 μ m. B) Box plot representations of the 400/480 nm fluorescence response ratios of roGFP2 at 6 days post-transfection. The results are normalized to the average value of the 'p15' condition, which is set to one arbitrary unit (a.u.). The bottom and top of each box represent the 25th and 75th percentile values, respectively; the horizontal line inside each box represents the median; and the horizontal lines below and above each box denote the mean minus and plus one standard deviation, respectively. The number of randomly chosen measurements (from at least 6 different cells) is indicated below each box plot. The data were statistically compared. C) Distribution pattern of roGFP2-PTS1 at 3 days post-transfection: PO (black boxes), C (white boxes) or PO/C (grey boxes). The number of cells analyzed is indicated above each condition.




Figure 2: The cysteine at position 11 of human PEX5 mediates the oxidative stress-related decline in PTS1 import efficiency. Primary (P; passage number: 24) and SV40 T-antigen-transformed (T) PEX5 null HuFs were transiently transfected with a plasmid encoding peroxisomal (roGFP2-PTS1) or cytosolic (roGFP2) roGFP2, in combination or not with a plasmid encoding HsPEX5WT (5_{WT}) or HsPEX5C11K (5_{C11K}). The cells were cultivated in the standard growth medium and analyzed 3 days post-transfection. A) Subcellular localizations of roGFP2-PTS1: peroxisomes (PO, black boxes), cytosol (C, white boxes) or peroxisomes/cytosol (PO/C, grey boxes); 100 randomly selected cells were analyzed per condition. B) Box plot representations of the 400/480 nm fluorescence response ratios of roGFP2. The results are normalized to the average value of the corresponding late-passage primary cells, which is set to one arbitrary unit (a.u.). The number of randomly chosen

measurements (from at least 10 different cells) is indicated below each box plot. For a description of these plots, see legend of Figure 1. The data were statistically compared.




Figure 3: The efficiency of PTS1 protein import complementation in aged primary PEX5 null human fibroblasts is affected by the redox state of the cytosol and the identity of the ubiquitinatable residue near the N-terminus of human PEX5. Primary PEX5 null HuFs (passage number: 25) were transiently transfected with a monocistronic plasmid encoding cytosolic roGFP2 (roGFP2) or a bicistronic plasmid encoding peroxisomal roGFP2 (roGFP2-PTS1) and wild-type HsPEX5 (5wt) or HsPEX5C11K (5C11K). After 3 days, the standard growth medium was replaced by fresh medium supplemented (+NAC) or not (-NAC) with 1 mM NAC. The cells were analyzed at the indicated time points. A) Box plot representations of the 400/480 nm fluorescence response ratios of roGFP2. The results are normalized to the average value of the '0 h' condition, which is set to one arbitrary unit (a.u.). The number of randomly chosen measurements (from at least 15 individual cells) is indicated below each box plot. For a description of these plots, see legend of Figure 1B. B and C) Subcellular localization of roGFP2-PTS1 under different treatment conditions (plotted as a function of time). PO, peroxisomal localization; PO/C, mixed peroxisomal and cytosolic localization.




Figure 4: Oxidized glutathione interferes with monoubiquitination of human PEX5 at Cys11. ³⁵S-labeled PEX5_{WT} and PEX5_{C11K} were subjected to in vitro import reactions (supplemented with GST-Ub) in the absence or presence of increasing concentrations of oxidized glutathione (GSSG) (for more details, see Materials and Methods). After 20 min, the samples were treated with NEM, and the organelles were isolated by centrifugation. The organelle fractions were subjected to SDS-PAGE under non-reducing (-DTT) and reducing (+DTT) conditions, and blotted onto a nitrocellulose membrane. The membrane was exposed to an X-ray film. Lane I, 10% of the ³⁵S-labeled PEX5 reticulocyte lysate used in each lane. The numbers to the left indicate the migration of the molecular mass markers (in kDa).